



A LONG JOURNEY OF GRAVITATIONAL WAVES



Universität Zürich UZH

Gravitation and Astrophysics Group

Y.Xu, E.Hamilton, D.Lopez, M.Ebersold, S.Tiwari, P.Jetzer

A long time ago in a galaxy far, far away ...


Strong gravitational waves can be produced by catastrophic events, such as colliding black holes and neutron stars, pulsars, supernovae, or gravitational radiation left over from the Big Bang.

COALESCING COMPACT BINARIES



Credit: Dana Berry, SkyWorks Digital, Inc.

ROTATING STARS (PULSARS)




Credit: NASA/Goddard Space Flight Center CIL

SUPERNOVA EXPLOSIONS



Credit: NASA, ESA, J. Hester and A. Löll

STOCHASTIC BACKGROUND




Credit: ESA and the Planck Collaboration - D. Ducros

Gravitational wave

Black hole

Spacetime

Gravitational waves are ripples in spacetime, traveling at the speed of light. They will disturb the spacetime thus any object in the path will get squeezed and stretched as the waves pass by.



When gravitational waves reach us from billions of light years away, they distort spacetime by an almost negligible amount. This distortion is many times smaller than the size of a proton nucleus. However, with the detectors targeting different frequency bands, they can still be measured...

Credit: ©Johan Jarnestad




1000 Hz - 10 Hz
Ground Based Detector

100 mHz - 0.1 mHz
Space Detector


320 nanoHz - 1 nanoHz
Pulsar Timing Array

10⁻¹³ Hz - 10⁻¹⁶ Hz
Microwave Background


Current and future ground based observatories such as LIGO, Virgo, and Einstein Telescope can detect longer wave-lengths than the detectors' lengths (a few kilometres), corresponding to periods of a few hundredths to a few thousandths of a second.




LISA, a trio of probes scheduled to launch in 2034, will have virtual arms millions of kilometres long, making it sensitive to waves with periods of a few seconds up to several hours. This will allow us to detect gravitational waves from supermassive black holes.



Gravitational waves from distant galaxies perturb the distance between Earth and stars in the Milky Way. The goal is to detect waves of periods lasting years by examining delays in the radio signals from spinning neutron stars known as pulsars.




The Universe's oldest measurable radiation could carry evidence of gravitational waves from the Big Bang. Those waves would not be detectable directly. However, their stretch across a significant fraction of the observable Universe could be detectable.




Credit: Davide Castelvecchi

Global Gravitational Waves Detector Network



Gravitational Waves Detections: Masses in the Stellar Graveyard



LIGO-Virgo-KAGRA | Aaron Geller | Northwestern

"Sounds" from space

